

**Clinical Policy: Trastuzumab/Biosimilars, Trastuzumab-Hyaluronidase**

Reference Number: CP.PHAR.228

Effective Date: 06.01.16

Last Review Date: 05.25

Line of Business: Commercial, HIM, Medicaid

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

**Description**

- Trastuzumab (Herceptin®) is a human epidermal growth factor receptor 2 (HER2)/neu receptor antagonist.
- Trastuzumab-dkst (Ogivri®), trastuzumab-pkrb (Herzuma®), trastuzumab-dttb (Ontruzant®), trastuzumab-qyyp (Trazimera®), trastuzumab-anns (Kanjinti®), and trastuzumab-strf (Hercessi™) are Herceptin biosimilars.
- Trastuzumab-hyaluronidase-oysk (Herceptin Hylecta™) is a combination of trastuzumab and hyaluronidase, an endoglycosidase.

**FDA Approved Indication(s)**

| Indications*             | Description   | Herceptin, Hercessi, Herzuma, Ogivri, Ontruzant, Trazimera, Kanjinti   | Herceptin Hylecta |
|--------------------------|---|--|-------------------|
| Adjuvant breast cancer   | For adjuvant treatment of HER2-overexpressing node positive or node negative (ER/PR negative or with one high risk feature**) breast cancer:                | As part of a treatment regimen consisting of doxorubicin, cyclophosphamide, and either paclitaxel or docetaxel | X                 |
|                          |   | As part of a treatment regimen with docetaxel and carboplatin  | X                 |
|                          |   | As a single agent following multi-modality anthracycline based therapy   | X                 |
| Metastatic breast cancer | In combination with paclitaxel for first-line treatment of HER2-overexpressing metastatic breast cancer   | X  | X                 |
|                          | As a single agent for treatment of HER2-overexpressing breast cancer in patients who have received one or more chemotherapy regimens for metastatic disease | X  | X                 |

**CLINICAL POLICY**

**Trastuzumab/Biosimilars, Trastuzumab-Hyaluronidase**

| Indications*   | Description   | Herceptin, Hercessi, Herzuma, Ogivri, Ontruzant, Trazimera, Kanjinti | Herceptin Hylecta |
|----------------|---|--|-------------------|
| Gastric cancer | In combination with cisplatin and capecitabine or 5-fluorouracil for the treatment of patients with HER2-overexpressing metastatic gastric or gastroesophageal junction (esophagogastric junction; EGJ) adenocarcinoma who have not received prior treatment for metastatic disease | X  | —                 |

\*Select patients for therapy based on an FDA-approved companion diagnostic for trastuzumab.

\*\* High-risk is defined as ER/PR positive with one of the following features: tumor size > 2 cm, age < 35 years, or tumor grade 2 or 3>

**Policy/Criteria**

*Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.*

It is the policy of health plans affiliated with Centene Corporation<sup>®</sup> that Herceptin/biosimilars and Herceptin Hylecta are **medically necessary** when the following criteria are met:

**I. Initial Approval Criteria**

**A. Breast Cancer** (must meet all):

1. Diagnosis of HER2-positive breast cancer or leptomeningeal metastases from HER2-positive breast cancer;
2. Prescribed by or in consultation with an oncologist;
3. Age ≥ 18 years;
4. If request is for Herceptin, Hercessi, Herzuma, or Ontruzant, member meets one of the following (a, b, or c):
  - a. If request if for Herceptin, member must use ALL of the following, unless clinically significant adverse effects are experienced or all are contraindicated (i and ii):
    - i. Kanjinti, Ogivri, Trazimera;
    - ii. If member has failed Kanjinti, Ogivri, and Trazimera, then member must use Ontruzant, Hercessi, and Herzuma;  
*\*Prior authorization may be required*
  - b. If request is for Herzuma, Hercessi, or Ontruzant, member must use ALL of the following, unless clinically significant adverse effects are experienced or all are contraindicated: Kanjinti, Ogivri, Trazimera;  
*\*Prior authorization may be required*
  - c. Request is for treatment associated with cancer for a State with regulations against step therapy in certain oncology settings (*see Appendix E*);

**CLINICAL POLICY**

**Trastuzumab/Biosimilars, Trastuzumab-Hyaluronidase**

5. Request meets one of the following (a, b, c, or d):\*
  - a. Herceptin, Ogivri, Herzuma, Hercessi, Ontruzant, Trazimera, Kanjinti: Dose does not exceed 8 mg/kg IV for adjuvant therapy or 4 mg/kg IV for treatment of metastatic disease (*see Appendix D for dose rounding guidelines*);
  - b. Herceptin, Ogivri, Herzuma, Hercessi, Ontruzant, Trazimera, Kanjinti: Intrathecal administration for leptomeningeal metastasis;
  - c. Herceptin Hylecta: Dose does not exceed 600 mg/10,000 units SC every 3 weeks (*see Appendix D for dose rounding guidelines*);
  - d. Dose/product is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*\*Prescribed regimen must be FDA-approved or recommended by NCCN*

**Approval duration: 6 months**

**B. Gastric, Esophageal and Esophagogastric Junction Cancer (must meet all):**

1. Diagnosis of HER2-positive gastric, esophageal, or EGJ adenocarcinoma;
2. Prescribed by or in consultation with an oncologist;
3. Age  $\geq$  18 years;
4. Disease is advanced, recurrent, unresectable, or metastatic;
5. Prescribed in combination with systemic chemotherapy;

*\*Prior authorization may be required.*

6. If request is for Herceptin, Hercessi, Herzuma, or Ontruzant, member meets one of the following (a, b, or c):
  - a. If request is for Herceptin, member must use ALL of the following, unless clinically significant adverse effects are experienced or all are contraindicated (i and ii):
    - i. Kanjinti, Ogivri, Trazimera;
    - ii. If member has failed Kanjinti, Ogivri, and Trazimera, then member must use Ontruzant, Hercessi, and Herzuma;

*\*Prior authorization may be required*

- b. If request is for Herzuma, Hercessi, or Ontruzant, member must use ALL of the following, unless clinically significant adverse effects are experienced or all are contraindicated: Kanjinti, Ogivri, Trazimera;

*\*Prior authorization may be required*

- c. Request is for treatment associated with cancer for a State with regulations against step therapy in certain oncology settings (*see Appendix E*);

6. Request meets one of the following (a or b):\*
  - a. Herceptin, Herzuma, Hercessi, Ogivri, Ontruzant, Trazimera, Kanjinti: Dose does not exceed 8 mg/kg IV (*see Appendix D for dose rounding guidelines*);
  - b. Dose/product is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*\*Prescribed regimen must be FDA-approved or recommended by NCCN*

**Approval duration: 6 months**

**C. Endometrial Carcinoma (off-label) (must meet all):**

1. Diagnosis of HER2-positive endometrial carcinoma with serous histology;
2. Prescribed by or in consultation with an oncologist;

**CLINICAL POLICY**

**Trastuzumab/Biosimilars, Trastuzumab-Hyaluronidase**

3. Age  $\geq$  18 years;
4. Disease is advanced (i.e., stage III/IV) or recurrent;
5. Prescribed in one of the following ways (a or b):
  - a. In combination with carboplatin and paclitaxel;\*  
*\*Prior authorization may be required.*
  - b. As a single agent for maintenance therapy;
6. If request is for Herceptin, Hercessi, Herzuma, or Ontruzant, member meets one of the following (a, b, or c):
  - a. If request is for Herceptin, member must use ALL of the following, unless clinically significant adverse effects are experienced or all are contraindicated (i and ii):
    - i. Kanjinti, Ogivri, Trazimera;
    - ii. If member has failed Kanjinti, Ogivri, and Trazimera, then member must use Ontruzant, Hercessi, and Herzuma;  
*\*Prior authorization may be required*
  - b. If request is for Herzuma, Hercessi, or Ontruzant, member must use ALL of the following, unless clinically significant adverse effects are experienced or all are contraindicated: Kanjinti, Ogivri, Trazimera;  
*\*Prior authorization may be required*
  - c. Request is for treatment associated with cancer for a State with regulations against step therapy in certain oncology settings (*see Appendix E*);
7. Dose is within FDA maximum limit for any FDA-approved indication or is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).\*  
*\*Prescribed regimen must be FDA-approved or recommended by NCCN*

**Approval duration: 6 months**

**D. Colorectal Cancer (off-label) (must meet all):**

1. Diagnosis of advanced or metastatic colorectal cancer and disease is all of the following (a, b, and c):
  - a. HER2 positive;
  - b. Wild-type *RAS* (defined as wild-type in both KRAS and NRAS [i.e., KRAS and NRAS mutation-negative] as determined by an FDA-approved test for this use);
  - c. Wild-type *BRAF* (i.e., BRAF mutation-negative);
2. Prescribed by or in consultation with an oncologist;
3. Age  $\geq$  18 years;
4. If request is for Herceptin, Hercessi, Herzuma, or Ontruzant, member meets one of the following (a, b, or c):
  - a. If request is for Herceptin, member must use ALL of the following, unless clinically significant adverse effects are experienced or all are contraindicated (i and ii):
    - i. Kanjinti, Ogivri, Trazimera;
    - ii. If member has failed Kanjinti, Ogivri, and Trazimera, then member must use Ontruzant, Hercessi, and Herzuma;  
*\*Prior authorization may be required*

**CLINICAL POLICY**

**Trastuzumab/Biosimilars, Trastuzumab-Hyaluronidase**

- b. If request is for Herzuma, Hercessi, or Ontruzant, member must use ALL of the following, unless clinically significant adverse effects are experienced or all are contraindicated: Kanjinti, Ogivri, Trazimera;  
*\*Prior authorization may be required*
- c. Request is for treatment associated with cancer for a State with regulations against step therapy in certain oncology settings(*see Appendix E*);
- 5. Prescribed in combination with Perjeta<sup>®</sup> (pertuzumab), Tykerb<sup>®</sup> (lapatinib), or Tukysa<sup>®</sup> (tucatinib);\*  
*\*Prior authorization may be required.*
- 6. Dose is within FDA maximum limit for any FDA-approved indication or is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).\*  
*\*Prescribed regimen must be FDA-approved or recommended by NCCN*

**Approval duration: 6 months**

**E. Salivary Gland Tumor (off-label) (must meet all):**

- 1. Diagnosis of HER2-positive salivary gland tumor;
- 2. Prescribed by or in consultation with an oncologist;
- 3. Age  $\geq$  18 years;
- 4. Disease is recurrent;
- 5. Prescribed in one of the following manners (a, b, or c):
  - a. Single agent;
  - b. Combination with docetaxel;\*
  - c. Combination with Perjeta;\**\*Prior authorization may be required.*
- 6. If request is for Herceptin, Hercessi, Herzuma, or Ontruzant, member meets one of the following (a, b, or c):
  - a. If request if for Herceptin, member must use ALL of the following, unless clinically significant adverse effects are experienced or all are contraindicated (i and ii):
    - i. Kanjinti, Ogivri, Trazimera;
    - ii. If member has failed Kanjinti, Ogivri, and Trazimera, then member must use Ontruzant, Hercessi, and Herzuma;  
*\*Prior authorization may be required*
  - b. If request is for Herzuma, Hercessi, or Ontruzant, member must use ALL of the following, unless clinically significant adverse effects are experienced or all are contraindicated: Kanjinti, Ogivri, Trazimera;  
*\*Prior authorization may be required*
  - c. Request is for treatment associated with cancer for a State with regulations against step therapy in certain oncology settings (*see Appendix E*);
- 7. Dose is within FDA maximum limit for any FDA-approved indication or is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).\*  
*\*Prescribed regimen must be FDA-approved or recommended by NCCN*

**Approval duration: 6 months**

**CLINICAL POLICY****Trastuzumab/Biosimilars, Trastuzumab-Hyaluronidase****F. Gallbladder Cancer or Cholangiocarcinoma (off-label) (must meet all):**

1. Diagnosis of HER2-positive gallbladder cancer or cholangiocarcinoma;
2. Prescribed by or in consultation with an oncologist;
3. Age  $\geq$  18 years;
4. Disease is unresectable, resected gross residual (R2), or metastatic;
5. Prescribed in combination with Perjeta or Tukysa\*;  
*\*Prior authorization may be required.*
6. If request is for Herceptin, Hercessi, Herzuma, or Ontruzant, member meets one of the following (a, b, or c):
  - a. If request is for Herceptin, member must use ALL of the following, unless clinically significant adverse effects are experienced or all are contraindicated (i and ii):
    - i. Kanjinti, Ogivri, Trazimera;
    - ii. If member has failed Kanjinti, Ogivri, and Trazimera, then member must use Ontruzant, Hercessi, and Herzuma;  
*\*Prior authorization may be required*
  - b. If request is for Herzuma, Hercessi, or Ontruzant, member must use ALL of the following, unless clinically significant adverse effects are experienced or all are contraindicated: Kanjinti, Ogivri, Trazimera;  
*\*Prior authorization may be required*
  - c. Request is for treatment associated with cancer for a State with regulations against step therapy in certain oncology settings (*see Appendix E*);
7. Dose is within FDA maximum limit for any FDA-approved indication or is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).\*  
*\*Prescribed regimen must be FDA-approved or recommended by NCCN*

**Approval duration: 6 months****G. Other diagnoses/indications (must meet all):**

1. One of the following (a, b, or c):
  - a. Request is for treatment associated with cancer for a State with regulations against step therapy in certain oncology settings (*see Appendix E*);
  - b. If request is for Herceptin, member must use ALL of the following, unless clinically significant adverse effects are experienced or all are contraindicated (i and ii):
    - i. Kanjinti, Ogivri, Trazimera;
    - ii. If member has failed Kanjinti, Ogivri, and Trazimera, then member must use Ontruzant, Hercessi, and Herzuma;  
*\*Prior authorization may be required*
  - c. If request is for Herzuma, Hercessi, or Ontruzant, member must use ALL of the following, unless clinically significant adverse effects are experienced or all are contraindicated: Kanjinti, Ogivri, Trazimera;  
*\*Prior authorization may be required*

## CLINICAL POLICY

### Trastuzumab/Biosimilars, Trastuzumab-Hyaluronidase

2. One of the following (a or b):
  - a. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (i or ii):
    - i. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace, and CP.PMN.255 for Medicaid; or
    - ii. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace, and CP.PMN.16 for Medicaid; or
  - b. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 2a above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

## II. Continued Approval

### A. All Indications in Section I (must meet all):

1. Currently receiving medication via Centene benefit, or documentation supports that member is currently receiving the requested agent for a covered indication and has received this medication for at least 30 days;
2. Member is responding positively to therapy;
3. For adjuvant breast cancer therapy, member has received  $\leq 52$  weeks of therapy total;
4. If request is for Herceptin, Hercessi, Herzuma, or Ontruzant, member meets one of the following (a, b, or c):
  - a. If request is for Herceptin, member must use ALL of the following, unless clinically significant adverse effects are experienced or all are contraindicated (I and ii):
    - i. Kanjinti, Ogivri, Trazimera;
    - ii. If member has failed Kanjinti, Ogivri, and Trazimera, then member must use Ontruzant, Hercessi, and Herzuma;  
*\*Prior authorization may be required*
  - b. If request is for Herzuma, Hercessi, or Ontruzant, member must use ALL of the following, unless clinically significant adverse effects are experienced or all are contraindicated: Kanjinti, Ogivri, Trazimera;  
*\*Prior authorization may be required*
  - c. Request is for treatment associated with cancer for a State with regulations against step therapy in certain oncology settings (*see Appendix E*);
5. If request is for a dose increase, request meets one of the following (a, b, or c):\*
  - a. Breast cancer (i, ii, or iii):
    - i. Herceptin, Ogivri, Hercessi, Herzuma, Ontruzant, Trazimera, Kanjinti: New dose does not exceed 8 mg/kg IV for adjuvant therapy or 4 mg/kg IV for

**CLINICAL POLICY**

**Trastuzumab/Biosimilars, Trastuzumab-Hyaluronidase**

- treatment of metastatic disease (*see Appendix D for dose rounding guidelines*);
- ii. Herceptin, Ogivri, Hercessi, Herzuma, Ontruzant, Trazimera, Kanjinti: Intrathecal administration for leptomeningeal metastases;
- iii. Herceptin Hylecta: New dose does not exceed 600 mg/10,000 units SC every 3 weeks (*see Appendix D for dose rounding guidelines*);
- b. Gastric, esophageal, EGJ cancer: Herceptin, Hercessi, Herzuma, Ogivri, Ontruzant, Trazimera, Kanjinti: New dose does not exceed 8 mg/kg IV (*see Appendix D for dose rounding guidelines*);
- c. New dose/product is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).  
*\*Prescribed regimen must be FDA-approved or recommended by NCCN*

**Approval duration: 12 months (total of 52 weeks for adjuvant breast cancer therapy)**

**B. Other diagnoses/indications (must meet all):**

1. One of the following (a, b, or c):
  - a. Request is for treatment associated with cancer for a State with regulations against step therapy in certain oncology settings (*see Appendix E*);
  - b. If request is for Herceptin, member must use ALL of the following, unless clinically significant adverse effects are experienced or all are contraindicated (i and ii):
    - i. Kanjinti, Ogivri, Trazimera;
    - ii. If member has failed Kanjinti, Ogivri, and Trazimera, then member must use Ontruzant, Hercessi, and Herzuma;  
*\*Prior authorization may be required*
  - c. If request is for Herzuma, Hercessi, or Ontruzant, member must use ALL of the following, unless clinically significant adverse effects are experienced or all are contraindicated: Kanjinti, Ogivri, Trazimera;  
*\*Prior authorization may be required*
2. One of the following (a or b):
  - a. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (i or ii):
    - i. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace, and CP.PMN.255 for Medicaid; or
    - ii. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace, and CP.PMN.16 for Medicaid; or
  - b. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 2a above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.



**CLINICAL POLICY**

**Trastuzumab/Biosimilars, Trastuzumab-Hyaluronidase**

**III. Diagnoses/Indications for which coverage is NOT authorized:**

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid, or evidence of coverage documents.

**IV. Appendices/General Information**

*Appendix A: Abbreviation/Acronym Key*

|  |  |
|--|--|
| BRAF: v-Raf murine sarcoma viral oncogene homolog B1 | KRAS: Kirsten rat sarcoma 2 viral oncogene homologue |
| FDA: Food and Drug Administration                    | NRAS: neuroblastoma RAS viral oncogene homologue     |
| EGJ: esophagogastric junction                        |  |
| HER2: human epidermal growth factor receptor 2       |  |

*Appendix B: Therapeutic Alternatives*

Not applicable

*Appendix C: Contraindications/Boxed Warnings*

- Contraindication(s): none reported
- Boxed warning(s): cardiomyopathy, embryo-fetal toxicity, pulmonary toxicity
  - Additionally, Herceptin, Ogivri, Herzuma, Ontruzant, Trazimera, Kanjinti, Hercessi for infusion reactions

*Appendix D: Dose Rounding Guidelines*

| Weight-based Dose Range   | Vial Quantity Recommendation           |
|---------------------------|--|
| ≤ 157.49 mg               | 1 vial of 150 mg                       |
| 157.5 mg to 314.99 mg     | 2 vials of 150 mg                      |
| 315 mg to 440.99 mg       | 1 vial of 420 mg                       |
| 441 mg to 598.49 mg       | 1 vial of 150 mg and 1 vial 420 mg     |
| 598.5 mg to 881.99 mg     | 2 vials of 420 mg                      |
| 882 mg to 1,039.49 mg     | 1 vial of 150 mg and 2 vials of 420 mg |
| 1,039.5 mg to 1,322.99 mg | 3 vials of 420 mg                      |

*Appendix E: States with Regulations against Redirections in Cancer*

| State | Step Therapy Prohibited? | Notes   |
|-------|--------------------------|---|
| FL    | Yes                      | For stage 4 metastatic cancer and associated conditions.  |
| GA    | Yes                      | For stage 4 metastatic cancer. Redirection does not refer to review of medical necessity or clinical appropriateness.     |
| IA    | Yes                      | For standard of care stage 4 cancer drug use, supported by peer-reviewed, evidence-based literature, and approved by FDA. |

**CLINICAL POLICY**

**Trastuzumab/Biosimilars, Trastuzumab-Hyaluronidase**

| State | Step Therapy Prohibited? | Notes   |
|-------|--------------------------|---|
| LA    | Yes                      | For stage 4 advanced, metastatic cancer or associated conditions. Exception if “clinically equivalent therapy, contains identical active ingredient(s), and proven to have same efficacy. |
| MS    | Yes                      | <i>*Applies to HIM requests only*</i><br>For advanced metastatic cancer and associated conditions   |
| NV    | Yes                      | Stage 3 and stage 4 cancer patients for a prescription drug to treat the cancer or any symptom thereof of the covered person  |
| OH    | Yes                      | <i>*Applies to Commercial and HIM requests only*</i><br>For stage 4 metastatic cancer and associated conditions   |
| OK    | Yes                      | <i>*Applies to HIM requests only*</i><br>For advanced metastatic cancer and associated conditions   |
| PA    | Yes                      | For stage 4 advanced, metastatic cancer   |
| TN    | Yes                      | For advanced metastatic cancer and associated conditions  |
| TX    | Yes                      | For stage 4 advanced, metastatic cancer and associated conditions   |

*Appendix F: General Information*

| Residual Tumor (R) Classification |                            |                                  |
|-----------------------------------|----------------------------|----------------------------------|
| R0                                | no residual tumor          | resected, negative margin        |
| R1                                | microscopic residual tumor | resected, positive margin        |
| R2                                | macroscopic residual tumor | resected, gross residual disease |

**V. Dosage and Administration**

| Drug Name   | Indication                        | Dosing Regimen   | Maximum Dose |
|---|-----------------------------------|--|--------------|
| Trastuzumab (Herceptin), Trastuzumab-dkst (Ogivri), Trastuzumab-dttb (Ontruzant), Trastuzumab-pkrb (Herzuma), Trastuzumab-qyyp (Trazimera), Trastuzumab-hyaluronidase -oysk | Adjuvant treatment, breast cancer | Administer according to one of the following doses and schedules for a total of 52 weeks:<br><b><u>Herceptin, Ogivri, Herzuma, Ontruzant, Trazimera, Kanjinti, Hercessi:</u></b><br>During and following paclitaxel, docetaxel, or docetaxel/carboplatin: <ul style="list-style-type: none"> <li>Initial dose of 4 mg/kg as an IV infusion over 90 minutes then at 2 mg/kg as an IV infusion over 30 minutes weekly during chemotherapy for the first 12 weeks (paclitaxel or docetaxel) or 18 weeks (docetaxel/carboplatin).</li> <li>One week following the last weekly dose of the trastuzumab product, administer trastuzumab product at 6 mg/kg as an IV infusion over 30 to 90 minutes every 3 weeks.</li> </ul> | 8 mg/kg      |

**CLINICAL POLICY**

**Trastuzumab/Biosimilars, Trastuzumab-Hyaluronidase**

| Drug Name   | Indication                          | Dosing Regimen   | Maximum Dose                      |
|---|-------------------------------------|--|-----------------------------------|
| (Herceptin Hylecta),<br>Trastuzumab-anns<br>(Kanjinti),<br>Trastuzumab-strf (Hercessi)  |                                     | <p><b><u>Herceptin, Ogivri, Herzuma, Ontruzant, Trazimera, Kanjinti, Hercessi:</u></b><br/>           As a single agent within 3 weeks following completion of multi-modality, anthracycline based chemotherapy regimens:</p> <ul style="list-style-type: none"> <li>• Initial dose: 8 mg/kg as an IV infusion over 90 minutes.</li> <li>• Subsequent doses: 6 mg/kg as an IV infusion over 30 to 90 minutes every 3 weeks</li> </ul>                                |                                   |
|   |                                     | <p><b><u>Herceptin Hylecta (subcutaneous product):</u></b><br/>           As part of a treatment regimen consisting of doxorubicin, cyclophosphamide, and either paclitaxel or docetaxel; as part of a treatment regimen with docetaxel and carboplatin; as a single agent following multi-modality anthracycline based therapy: 600 mg trastuzumab and 10,000 units hyaluronidase administered subcutaneously over approximately 2-5 minutes once every 3 weeks</p> | 600 mg/10,000 units every 3 weeks |
| Trastuzumab (Herceptin),<br>Trastuzumab-dkst (Ogivri),<br>Trastuzumab-dttb (Ontruzant),<br>Trastuzumab-pkrb (Herzuma),<br>Trastuzumab-qyyp (Trazimera),<br>Trastuzumab-hyaluronidase -oysk (Herceptin Hylecta),<br>Trastuzumab-anns (Kanjinti), | Metastatic treatment, breast cancer | <p><b><u>Herceptin, Ogivri, Herzuma, Ontruzant, Trazimera, Kanjinti, Hercessi:</u></b><br/>           As a single agent, or in combination with paclitaxel, at an initial dose of 4 mg/kg as a 90-minute intravenous infusion followed by subsequent once weekly doses of 2 mg/kg as 30-minute intravenous infusions until disease progression.</p>  | 4 mg/kg                           |
|   |                                     | <p><b><u>Herceptin Hylecta (subcutaneous product):</u></b><br/>           As a single agent or in combination with paclitaxel: 600 mg trastuzumab and 10,000 units hyaluronidase administered subcutaneously over approximately 2-5 minutes once every 3 weeks.</p>  | 600 mg/10,000 units every 3 weeks |

**CLINICAL POLICY**

**Trastuzumab/Biosimilars, Trastuzumab-Hyaluronidase**

| Drug Name   | Indication                | Dosing Regimen  | Maximum Dose |
|---|---------------------------|---|--------------|
| Trastuzumab-strf (Hercessi)   |                           |   |              |
| Trastuzumab (Herceptin),<br>Trastuzumab-dkst (Ogivri),<br>Trastuzumab-dttb (Ontruzant),<br>Trastuzumab-qyyp (Trazimera),<br>Trastuzumab-anns (Kanjinti),<br>Trastuzumab-strf (Hercessi) | Metastatic gastric cancer | <b><u>Herceptin, Herzuma, Ogivri, Ontruzant, Trazimera, Kanjinti, Hercessi:</u></b><br>Administer at an initial dose of 8 mg/kg as a 90 minute intravenous infusion followed by subsequent doses of 6 mg/kg as an intravenous infusion over 30 to 90 minutes every three weeks until disease progression. | 8 mg/kg      |

**VI. Product Availability**

| Drug Name  | Availability*  |
|--|--|
| Trastuzumab (Herceptin)                            | Single-dose vial: 150 mg   |
| Trastuzumab-dkst (Ogivri)                          | Single-dose vial: 150 mg<br>Multi-dose vial: 420 mg**                    |
| Trastuzumab-pkrb (Herzuma)                         | Single-dose vial: 150 mg<br>Multi-dose vial: 420 mg                      |
| Trastuzumab-dttb (Ontruzant)                       | Single-dose vial: 150 mg<br>Multi-dose vial: 420 mg                      |
| Trastuzumab-qyyp (Trazimera)                       | Single-dose vial: 150 mg<br>Multi-dose vial: 420 mg                      |
| Trastuzumab-hyaluronidase-oysk (Herceptin Hylecta) | Single-dose vial: 600 mg (trastuzumab)/10,000 units (hyaluronidase)/5 mL |
| Trastuzumab-anns (Kanjinti)                        | Single-dose vial: 150 mg<br>Multi-dose vial: 420 mg                      |
| Trastuzumab-strf (Hercessi)                        | Single-dose vial: 150 mg<br>Multi-dose vial: 420 mg                      |

\*All products are supplied as a powder for reconstitution with the exception of Herceptin Hylecta which is supplied as a solution.

\*\* Product available with or without diluent provided

## CLINICAL POLICY

### Trastuzumab/Biosimilars, Trastuzumab-Hyaluronidase

#### VII. References

1. Herceptin Prescribing Information. South San Francisco, CA: Genentech, Inc.; June 2024. Available at: [https://www.gene.com/download/pdf/herceptin\\_prescribing.pdf](https://www.gene.com/download/pdf/herceptin_prescribing.pdf). Accessed January 13, 2025.
2. Ogivri Prescribing Information. Morgantown, WV: Mylan GmbH.; November 2024. Available at <https://www.ogivri.com/>. Accessed January 13, 2025.
3. Herzuma Prescribing Information. North Wales, PA: Teva Pharmaceuticals USA, Inc.; November 2024. <https://www.herzuma.com/>. Accessed January 13, 2025.
4. Ontruzant Prescribing Information. Jersey City, NJ: Organon; June 2021. <https://www.ontruzant.com/>. Accessed January 13, 2025.
5. Trazimera Prescribing Information. New York, NY: Pfizer Labs; November 2020. Available at <https://labeling.pfizer.com/ShowLabeling.aspx?id=12725>. Accessed January 13, 2025.
6. Herceptin Hylecta Prescribing Information. South San Francisco, CA: Genentech, Inc.; June 2024. Available at [https://www.gene.com/download/pdf/herceptin\\_hylecta\\_prescribing.pdf](https://www.gene.com/download/pdf/herceptin_hylecta_prescribing.pdf). Accessed January 13, 2025.
7. Kanjinti Prescribing Information. Thousand Oaks, CA: Amgen, Inc.; December 2024. Available at [https://www.pi.amgen.com/-/media/Project/Amgen/Repository/pi-amgen-com/Kanjinti/kanjinti\\_pi.pdf](https://www.pi.amgen.com/-/media/Project/Amgen/Repository/pi-amgen-com/Kanjinti/kanjinti_pi.pdf). Accessed January 13, 2025.
8. Hecessi Prescribing Information. Raleigh, NC: Accord BioPharma Inc.; September 2024. Available at: <https://dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=285d2142-5bfa-401c-9246-7de21c4451af>. Accessed January 13, 2025.
9. National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at: [http://www.nccn.org/professionals/drug\\_compendium](http://www.nccn.org/professionals/drug_compendium). Accessed January 27, 2025.
10. Fahrenbruch R, Kintzel P, Bott AM., et al. Dose rounding of biologic and cytotoxic anticancer agents: a position statement of the hematology/oncology pharmacy association. *Journal of Oncology Practice*. 2018;14(3)e130-e136.
11. National Comprehensive Cancer Network. Breast Cancer Version 6.2024. Available at: [https://www.nccn.org/professionals/physician\\_gls/pdf/breast.pdf](https://www.nccn.org/professionals/physician_gls/pdf/breast.pdf). Accessed January 27, 2025.
12. Hermanek P and Wittekind C. Residual tumor (R) classification and prognosis. *Semin Surg Oncol*. 1994 Jan-Feb;10(1):12-20

#### Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

| HCPCS Codes | Description   |
|-------------|---|
| J9355       | Injection, trastuzumab, excludes biosimilar, 10 mg          |
| J9356       | Injection, trastuzumab, 10 mg and hyaluronidase-oysk        |
| Q5112       | Injection, trastuzumab-dttb, biosimilar, (Ontruzant), 10 mg |
| Q5113       | Injection, trastuzumab-pkrb, biosimilar, (Herzuma), 10 mg   |
| Q5114       | Injection, trastuzumab-dkst, biosimilar, (Ogivri), 10 mg    |
| Q5116       | Injection, trastuzumab-qyyp, biosimilar, (Trazimera), 10 mg |
| Q5117       | Injection, trastuzumab-anns, biosimilar, (Kanjinti), 10 mg  |

**CLINICAL POLICY**

**Trastuzumab/Biosimilars, Trastuzumab-Hyaluronidase**

| <b>HCPCS Codes</b> | <b>Description</b>  |
|--------------------|---|
| Q5146              | Injection, trastuzumab-strf (hercessi), biosimilar, 10 mg |

| <b>Reviews, Revisions, and Approvals</b>   | <b>Date</b> | <b>P&amp;T Approval Date</b> |
|--|-------------|------------------------------|
| 2Q 2021 annual review: revised requirement of medical justification for inability to use preferred Ogivri or Trazimera to “must use” language and applied redirection to preferred biosimilars to other diagnoses/indications; added criteria for salivary gland tumor criteria for Herceptin as it is a NCCN-supported off-label indication; per NCCN support, added wild-type <i>BRAF</i> criterion for colorectal cancer and choice of oxaliplatin, in addition to cisplatin, for combination treatment of gastric cancers; updated product availability for Herceptin, Kanjinti, and Trazimera; references for HIM line of business off-label use revised from HIM.PHAR.21 to HIM.PA.154; references reviewed and updated. | 03.25.21    | 05.21                        |
| Per August SDC and prior clinical guidance, modified biosimilar redirection requirements for Herceptin to require use of Ogivri, Trazimera, Kanjinti, Ontruzant and Herzuma in a step-wise manner; for Ontruzant and Herzuma modified redirection to require use of Ogivri, Trazimera, and Kanjinti; for salivary gland tumor indication added redirection to preferred biosimilars per NCCN Compendium; adding legacy Wellcare Medicaid line of business (WCG.CP.PHAR.228 to be retired); added Nevada to Appendix E.   | 08.25.21    | 11.21                        |
| 2Q 2022 annual review: added qualifiers of “advanced” and “recurrent” for gastric, esophageal, or EGJ adenocarcinoma; initial approval durations were consolidated to 6 months for alignment between legacy WCG and other lines of business; removed general description of “stage IV or metastatic” cancer for states with regulations against redirections; clarified other diagnoses section to clarify intent for biosimilar steerage; references reviewed and updated.  | 02.16.22    | 05.22                        |
| Template changes applied to other diagnoses/indications.   | 10.10.22    |                              |
| 2Q 2023 annual review: added gallbladder cancer and cholangiocarcinoma as NCCN supported off-label indication; references reviewed and updated.  | 01.20.23    | 05.23                        |
| Updated Appendix E to include Oklahoma.  | 06.07.23    |                              |
| 2Q 2024 annual review: for adjuvant breast cancer continued therapy, added member has received ≤ 52 weeks of therapy per PI; for gastric, esophageal, or EGJ, added option for unresectable disease, revised prescribed combination therapy to “systemic chemotherapy” as additional regimens options available per NCCN; for endometrial carcinoma added option to be prescribed as single agent for maintenance therapy per NCCN; for colorectal cancer, removed   | 01.18.24    | 05.24                        |

**CLINICAL POLICY**

**Trastuzumab/Biosimilars, Trastuzumab-Hyaluronidase**

| Reviews, Revisions, and Approvals   | Date     | P&T Approval Date |
|---|----------|-------------------|
| requirement for no previous use of HER2 inhibitor therapy and added tucatinib as option to be prescribed in combination with; for gallbladder cancer or cholangiocarcinoma, added option for treatment with resected gross residual (R2) disease per NCCN; residual (R) tumor classification added to Appendix F; for Ogivri, updated product availability of 420 mg multi-dose vial supplied with or without diluent; references reviewed and updated. |          |                   |
| RT4: added Hercessi to policy as non-preferred biosimilar. Updated Appendix E to include Mississippi.   | 06.05.24 |                   |
| RT4: added new multi-dose vial formulation for Hercessi; HCPCS code added [Q5146] and removed codes [J3590, C9399]  | 11.19.24 |                   |
| 2Q 2025 annual review: for gallbladder cancer or cholangiocarcinoma, added option to be prescribed with tucatinib per NCCN; references reviewed and updated.  | 01.13.25 | 05.25             |

**Important Reminder**

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions, and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan

## CLINICAL POLICY

### Trastuzumab/Biosimilars, Trastuzumab-Hyaluronidase

retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment, or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members, and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

**Note: For Medicaid members**, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2016 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene<sup>®</sup> and Centene Corporation<sup>®</sup> are registered trademarks exclusively owned by Centene Corporation.